

**INFORMATION DISCLOSURE CITATION**

(Use several sheets if necessary)

PTO Form 1449

Attorney Docket 044508-5023

Application No. 10/528,800

Applicants: Christopher C. Broder *et al.*

Page 1 of 2

Filing Date: March 31, 2006

Group Art Unit: 1648

**OTHER DOCUMENTS (Including Author, Title, Date, Pertinent Pages, etc.)**

- |    |   |
|----|---|
| 1  | Bossart <i>et al.</i> (2001). Functional Expression and Membrane Fusion Tropism of The Envelope Glycoproteins of Hendra Virus. <i>Virology</i> 290:121-135.   |
| 2  | Chambers <i>et al.</i> (1990). Heptad repeat sequences are located adjacent to hydrophobic regions in several types of virus fusion glycoproteins. <i>J. Gen. Virol.</i> 71:3075-3080.  |
| 3  | Chua <i>et al.</i> (2000). Nipah Virus: A Recently Emergent Deadly Paramyxovirus. <i>Science</i> 288:1432-1435.   |
| 4  | Gosh <i>et al.</i> (1997). A Leucine Zipper Motif in the Ectodomain of Sendai Virus Fusion Protein Assembles in Solution and in Membranes and Specifically Binds Biologically-Active Peptides and the Virus. <i>Biochemistry</i> 36:15451-15462.    |
| 5  | Harcourt <i>et al.</i> (2001). Molecular Characterization of Nipah Virus, a Newly Emergent Paramyxovirus. <i>Virology</i> 271:334-349.  |
| 6  | Harcourt <i>et al.</i> (2001). Molecular Characterization of the Polymerase Gene and Genomic Termini of Nipah Virus. <i>Virology</i> 287:192-201.   |
| 7  | Joshi <i>et al.</i> (1998). A Core Trimer of the Paramyxovirus Fusion Protein: Parallels to Influenza Virus Hemagglutinin and HIV-1 gp41. <i>Virology</i> 248:20-34.  |
| 8  | Lambert <i>et al.</i> (1996). Peptides from conserved regions of paramyxovirus fusion (F) proteins are potent inhibitors of viral fusion. <i>Proc. Natl. Acad. Sci. US</i> 93:2186-2191.  |
| 9  | Paterson <i>et al.</i> (1987). Comparison of the Relative Roles of the F and HN Surface Glycoproteins of the Paramyxovirus Simian Virus 5 in Inducing Protective Immunity. <i>J. Virol.</i> 61:1972-1977.   |
| 10 | Rapaport <i>et al.</i> (1995). A synthetic peptide corresponding to a conserved heptad repeat domain is a potent inhibitor of Sendai virus-cell fusion: an emerging similarity with functional domains of other viruses. <i>EMBO</i> 14: 5524-5531. |
| 11 | Wang <i>et al.</i> (2001) Molecular biology of Hendra and Nipah Viruses. <i>Microbes and Infection</i> 3:279-287.   |
| 12 | Wild <i>et al.</i> (1997). Inhibition of measles virus infection and fusion with peptides corresponding to the leucine zipper region of the fusion protein. <i>J. Gen. Virol.</i> 78: 107-111.  |
| 13 | Yao <i>et al.</i> (1997). Association of the Parainfluenza Virus Fusion and Hemagglutinin-Neuraminidase Glycoproteins on Cell Surfaces. <i>J. Virol.</i> 71:650-656.  |
| 14 | Young <i>et al.</i> (1997). Analysis of Peptide Inhibitor of Paramyxovirus (NDV) Fusion Using Biological Assays, NMR, and Molecular Modeling. <i>Virology</i> 238: 291-304.   |
| 15 | Young <i>et al.</i> (1999). Interaction of Peptides with Sequences from the Newcastle Disease Virus Fusion Protein Heptad Repeat Regions. <i>J. Virol.</i> 73: 5954-5956.   |

Examiner

Date Considered

Examiner: Initial if reference considered, whether or not citation is in conformance with MPEP 609; draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.